

# King Philip Middle School

## Allergy Action Plan

Student's Name: \_\_\_\_\_ D.O.B: \_\_\_\_\_ Teacher: \_\_\_\_\_



ALLERGY TO: \_\_\_\_\_

Asthmatic Yes\*  No  \*Higher risk for severe reaction

### ◆ STEP 1: TREATMENT ◆

#### Symptoms:

- If a food allergen has been ingested, but *no symptoms*:
- Mouth Itching, tingling, or swelling of lips, tongue, mouth
- Skin Hives, itchy rash, swelling of the face or extremities
- Gut Nausea, abdominal cramps, vomiting, diarrhea
- Throat + Tightening of throat, hoarseness, hacking cough
- Lung + Shortness of breath, repetitive coughing, wheezing
- Heart + Thready pulse, low blood pressure, fainting, pale, blueness
- Other + \_\_\_\_\_
- If reaction is progressing (several of the above areas affected), give

#### Give Checked Medication\*\*:

- |                                 |  |
|---------------------------------|--|
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> EpiPen | <input type="checkbox"/> Antihistamine |

To be determined by physician authorizing treatment

The severity of symptoms can quickly change. + Potentially life-threatening.

#### DOSAGE

Epinephrine: inject intramuscularly (circle one) EpiPen EpiPen Jr. (see reverse side for instructions)

Antihistamine: give \_\_\_\_\_ medication/dose/route

Other: give \_\_\_\_\_ medication/dose/route

### ◆ STEP 2: EMERGENCY CALLS ◆

1. Call 911 (or Rescue Squad: \_\_\_\_\_). State that an allergic reaction has been treated, and additional epinephrine may be needed)

2. Dr. \_\_\_\_\_ at \_\_\_\_\_

3. Emergency contacts:

Name/Relationship	Phone Number(s)
a. _____	1.) _____ 2.) _____
b. _____	1.) _____ 2.) _____
c. _____	1.) _____ 2.) _____

**EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY!**

Parent/Guardian Signature \_\_\_\_\_

Date \_\_\_\_\_

Doctor's Signature \_\_\_\_\_ (Required)


Date \_\_\_\_\_

### EpiPen® and EpiPen® Jr. Directions

- Pull off gray activation cap.



- Hold black tip near outer thigh (always apply to thigh).



- Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds.

### Twinject™ 0.9 mg and Twinject™ 0.15 mg Directions



- Pull off green end cap, then red end cap.
- Put gray cap against outer thigh, press down firmly until needle penetrates. Hold for 10 seconds, then remove.



#### SECOND DOSE ADMINISTRATION:

If symptoms don't improve after 10 minutes, administer second dose:

- Unscrew gray cap and pull syringe from barrel by holding blue collar at needle base.
- Slide yellow or orange collar off plunger.
- Put needle into thigh through skin, push plunger down all the way, and remove.



DIRECTIONS FOR USE ARE ALSO ON EPIPEN